

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 May 2008 (15.05.2008)

PCT

(10) International Publication Number
WO 2008/055966 A1

(51) International Patent Classification:
A61K 9/20 (2006.01) *A61K 31/505* (2006.01)

Christian [DE/DE]; Meckenheimerstr. 20D, 67454 Has-
sloch (DE).

(21) International Application Number:
PCT/EP2007/062101

(74) Agent: REITSTÖTTER KINZEBACH & PARTNER;
Sternwartstr. 4, 81679 Munich (DE).

(22) International Filing Date:
8 November 2007 (08.11.2007)

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH,
CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG,
ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK,
LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW,
MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL,
PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY,
TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
06023367.3 9 November 2006 (09.11.2006) EP
60/999,579 19 October 2007 (19.10.2007) US

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL,
PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): ABBOTT
GMBH & CO. KG [DE/DE]; Max-Planck-Ring 2, 65205
Wiesbaden (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): LIEPOLD,
Bernd [DE/DE]; Birkenweg 44, 69221 Dossenheim
(DE). ROSENBERG, Jörg [DE/DE]; Bruchstrasse 29,
67156 Ellerstadt (DE). KNOBLOCH, Martin [DE/DE];
Medenheimerstr. 22a, 67141 Neuhofen (DE). NEHEN,

Published:

— with international search report

(54) Title: PHARMACEUTICAL DOSAGE FORM FOR ORAL ADMINISTRATION OF TYROSINE KINASE INHIBITOR

(57) Abstract: A pharmaceutical dosage form comprises a solid dispersion product of at least one tyrosine kinase inhibitor, at least one pharmaceutically acceptable polymer, and at least one pharmaceutically acceptable solubilizer.

WO 2008/055966 A1

Pharmaceutical dosage form for oral administration of tyrosine kinase inhibitor

5 The present invention relates to a pharmaceutical dosage form for oral administration of tyrosine kinase inhibitors, a method of preparing the dosage form and a method of treating proliferative disorders.

10 Tyrosine kinase inhibiting compounds are useful for treating diseases caused or exacerbated by upregulation or overexpression of protein tyrosine kinases. Unfortunately, the crystalline forms of many known tyrosine kinase inhibitors are characterized by a more or less pronounced poor solubility in aqueous liquids which affects their dissolution rate and bioavailability.

15 A measure of the potential usefulness of an oral dosage form of a pharmaceutical agent is the bioavailability observed after oral administration of the dosage form. Various factors can affect the bioavailability of a drug when administered orally. These factors include aqueous solubility, drug absorption throughout the gastrointestinal tract, dosage strength and first-pass effect. Aqueous solubility is one of the most important of these factors.

20 For a variety of reasons, such as patient compliance and taste masking, a solid dosage form is usually preferred over a liquid dosage form. In most instances, however, oral solid dosage forms of a drug provide a lower bioavailability than oral solutions of the drug.

25 There have been attempts to improve the bioavailability provided by solid dosage forms by forming solid solutions of the drug. Solid solutions are preferred physical systems because the components therein readily form liquid solutions when contacted with a liquid medium such as gastric juice. The ease of dissolution may be attributed at least in part to the fact that the energy required for dissolution of the components from a solid solution is less than that required for the dissolution of the components from a crystalline or microcrystalline solid phase. It is, however, important that the drug released from the solid solution remains water-solubilized in the aqueous fluids of the gastrointestinal tract; otherwise, the drug may precipitate in the gastrointestinal tract, resulting in low bioavailability.

35 WO 01/00175 discloses mechanically stable pharmaceutical dosage forms which are solid solutions of active ingredients in an auxiliary agent matrix. The matrix contains a homopolymer or a copolymer of N-vinyl pyrrolidone and a liquid or semi-solid surfactant.

40 WO 00/57854 discloses mechanically stable pharmaceutical dosage forms for peroral administration which contain at least one active compound, at least one thermo-

2

plastically mouldable, matrix-forming auxiliary and more than 10 and up to 40% by weight of a surface-active substance that has an HLB of between 2 and 18, is liquid at 20 °C, or has a drop point at between 20 and 50 °C.

- 5 US 2005/0208082 discloses a solubilizing composition comprising a mixture of vitamin E TPGS and linoleic acid. The solubilizing composition is used to disperse a lipophile in an aqueous phase. The lipophile may be a therapeutically effective lipophile such as lipophilic vitamins, coenzyme Q10, carotenoids, alpha-lipoic acid, essential fatty acids.
- 10 US 2005/0236236 discloses pharmaceutical compositions for administration of hydrophobic drugs, particularly steroids. The pharmaceutical compositions include a hydrophobic drug, a vitamin E substance and a surfactant. The reference claims a synergistic effect between the hydrophobic drug and the vitamin E substance.
- 15 There is a continuing need for the development of improved oral solid dosage forms of tyrosine kinase inhibitors.

The invention relates to a pharmaceutical dosage form which comprises a solid dispersion product of at least one tyrosine kinase inhibitor, at least one pharmaceutically acceptable polymer, and at least one pharmaceutically acceptable solubilizer.

In the dosage forms of the invention, the active ingredient is present as a solid dispersion or, preferably, as a solid solution. The term "solid dispersion" defines a system in a solid state (as opposed to a liquid or gaseous state) comprising at least two components, wherein one component is dispersed evenly throughout the other component or components. For example, the active ingredient or combination of active ingredients is dispersed in a matrix comprised of the pharmaceutically acceptable polymer(s) and pharmaceutically acceptable solubilizers. The term "solid dispersion" encompasses systems having small particles, typically of less than 1 µm in diameter, of one phase dispersed in another phase. When said dispersion of the components is such that the system is chemically and physically uniform or homogeneous throughout or consists of one phase (as defined in thermodynamics), such a solid dispersion will be called a "solid solution" or a "glassy solution". A glassy solution is a homogeneous, glassy system in which a solute is dissolved in a glassy solvent. Glassy solutions and solid solutions are preferred physical systems. These systems do not contain any significant amounts of active ingredients in their crystalline or microcrystalline state, as evidenced by thermal analysis (DSC) or X-ray diffraction analysis (WAXS).

The dosage forms according to the invention are characterized by an excellent stability and, in particular, exhibit high resistance against recrystallization or decomposition of the active ingredient(s).

The dosage forms of the present invention exhibit a release and absorption behaviour that is characterized by high attainable AUC (area under the plasma concentration-time

curve from 0 to 48 hours), high attainable C_{\max} (maximum plasma concentration), and low T_{\max} (time to reach maximum plasma concentration).

5 The term "AUC" means "Area Under the Curve" and is used in its normal meaning, i. e. as the area under the plasma concentration-time curve. " AUC_{0-48} " and " $AUC_{0-\infty}$ " refer to the area under the plasma concentration-time curve from 0 to 48 hours or from 0 hours to infinity, respectively.

10 In a preferred embodiment the invention provides a dosage form wherein said tyrosine kinase inhibitor is N-[4-(3-amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)-urea (ABT 869) (or a hydrate, solvate, N-oxide, or a pharmaceutically acceptable acid or base addition salt thereof). When administered to a human patient, in certain em-
bodiments, the dosage form produces a plasma profile characterized by a C_{\max} for ABT
15 869 from about 0.015 $\mu\text{g/mL/mg}$ to about 0.027 $\mu\text{g/mL/mg}$, in particular about 0.023 \pm 0.004 $\mu\text{g/mL/mg}$ (mean \pm SD), after a single dose.

When administered to the human patient, in certain embodiments, the dosage form produces a plasma profile characterized by a T_{\max} for ABT 869 of about 1 to about 3
20 hours, in particular about 2.8 \pm 0.6 hours, after a single dose.

In particular embodiments, when administered to the human patient, the dosage form produces an AUC_{0-48} per mg of ABT 869 from about 0.23 $\mu\text{g}\cdot\text{hr/mL/mg}$ to about 0.56
25 $\mu\text{g}\cdot\text{hr/mL/mg}$, in particular about 0.40 \pm 0.10 $\mu\text{g}\cdot\text{h/mL/mg}$, or an $AUC_{0-\infty}$ per mg of ABT 869 from about 0.27 $\mu\text{g}\cdot\text{hr/mL/mg}$ to about 0.81 $\mu\text{g}\cdot\text{hr/mL/mg}$, in particular about 0.55 \pm 0.17 $\mu\text{g}\cdot\text{h/mL/mg}$, per mg of dose after a single dose.

The plasma concentration profile may suitably be established in a group of at least ten healthy humans under fasting conditions, based on blood sampling at 0, 1, 3, 4, 6, 8,
30 24 and 48 hours. "Fasting conditions" means that the patients abstain from food or drink consumption except water and concomitant medications for 2 hours prior to and after dosing. Once the concentration-time points have been determined, the plasma concentration profile may be calculated, e. g. by a computer program or by the trape-
zoidal method. Administration of single dose of 10 mg ABT 869 to a human is consid-
35 ered suitable for determining the AUC values as used herein.

A preferred feature of the dosage form is their ability to release fine particles having, e. g., an average particle size of less than about 1000 nm, preferably less than about 800 nm, in particular less than about 500 nm and especially preferred less than about 200 nm, when the dosage form is brought into contact with an aqueous liquid. The fine
40 particles contain solubilised tyrosine kinase inhibitor, preferably in an essentially non-crystalline state. When the dosage form is administered orally, the aqueous liquid will be gastric juices. For in vitro testing purposes, the aqueous liquid may suitably be a volume of 900 ml of 1 N hydrochloric acid (USP apparatus II).

The dispersion formed upon contact with an aqueous liquid may also be useful as such, for example as oral liquid dosage form or parenteral injections.

Generally, the solid dispersion product comprises


- 5 from about 0.5 to 40 % by weight, preferably from about 1 to 25 % by weight, of said at least one tyrosine kinase inhibitor,
from about 40 to 97.5 % by weight, preferably from about 50 to 94 % by weight, of said at least one pharmaceutically acceptable polymer,
10 from about 2 to 20 % by weight, preferably from about 5 to 20 % by weight, of said at least one solubilizer, and
from about 0 to 15 % by weight, preferably from about 0 to 10 % by weight, of additives.

- Whereas the dosage form of the invention may consist entirely of solid dispersion
15 product, additives and adjuvants are usually used in formulating the solid dispersion product into the dosage forms. Generally, the dosage form comprises at least 10 % by weight, preferably at least 40 % by weight, and most preferred at least 45 % by weight, of solid dispersion product, based on the total weight of the solid dosage form.

- 20 Typically, a single dosage form of the invention contains the equivalent of about 0.1 mg to about 100 mg, preferably about 1.0 mg to about 50 mg, in particular 2.5 mg to 25 mg, of said at least one tyrosine kinase inhibitor.

- The inventive dosage form comprises a tyrosine kinase inhibitor or a combination of
25 two or more tyrosine kinase inhibitors. The dosage form may comprise a combination of one or more tyrosine kinase inhibitors and at least one further active ingredient. Various kinds of tyrosine kinase inhibitors can be effectively utilized.

- A preferred tyrosine kinase inhibitor is ABT 869 [N-[4-(3-amino-1H-indazol-4-yl)phenyl]-
30 N'-(2-fluoro-5-methylphenyl)urea] the preparation of which is described in WO 04/113304. The molecular structure of ABT 869 is depicted below:



35

A further preferred tyrosine kinase inhibitor is N-(4-(4-aminothieno[2,3-d]pyrimidin-5-yl)phenyl)-N'-(2-fluoro-5-(trifluoromethyl)phenyl)urea the preparation of which is described in US 2007/0155758.

5

Further tyrosine kinase inhibitors which may be used include sorafenib (trade name Nexavar), dasatinib, lapatinib (trade name Tykerb), imatinib (trade name Gleevec), motesanib, vandetanib (Zactima), MP-412, lestaurtinib, XL647, XL999, tandutinib, PKC412, nilotinib, AEE788, OSI-930, OSI-817, sunitinib maleate (trade name Sutent) and axitinib.

The term "tyrosine kinase inhibitors" is intended to encompass the hydrates, solvates (such as alcoholates), N-oxides, pharmaceutically acceptable acid or base addition salts of tyrosine kinase inhibiting compounds.

Pharmaceutically acceptable acid addition salts comprise the acid addition salt forms which can be obtained conveniently by treating the base form of the active ingredient with appropriate organic and inorganic acids.

Active ingredients containing an acidic proton may be converted into their non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases.

The invention is particularly useful for water-insoluble or poorly water-soluble (or "hydrophobic" or "lipophilic") compounds. Compounds are considered water-insoluble or poorly water-soluble when their solubility in water at 25 °C is less than 1 g/100 ml, especially less than 0,1 g/100 ml.

The term "pharmaceutically acceptable solubilizer" as used herein refers to a pharmaceutically acceptable non-ionic surfactant. The solubilizer may effectuate an instantaneous emulsification of the active ingredient released from the dosage form and/or prevent precipitation of the active ingredient in the aqueous fluids of the gastrointestinal tract. A single solubilizer as well as combinations of solubilizers may be used. According to an embodiment of the invention, the solid dispersion product comprises a combination of two or more pharmaceutically acceptable solubilizers.

Preferred solubilizers are selected from sorbitan fatty acid esters, polyalkoxylated fatty acid esters such as, for example, polyalkoxylated glycerides, polyalkoxylated sorbitan fatty acid esters or fatty acid esters of polyalkylene glycols, polyalkoxylated ethers of fatty alcohols, tocopheryl compounds or mixtures of two or more thereof. A fatty acid chain in these compounds ordinarily comprises from 8 to 22 carbon atoms. The polyalkylene oxide blocks comprise on average from 4 to 50 alkylene oxide units, preferably ethylene oxide units, per molecule.

Suitable sorbitan fatty acid esters are sorbitan monolaurate, sorbitan monopalmitate, sorbitan monostearate (Span® 60), sorbitan monooleate (Span® 80), sorbitan

tristearate, sorbitan trioleate, sorbitan monostearate, sorbitan monolaurate or sorbitan monooleate.


Examples of suitable polyalkoxylated sorbitan fatty acid esters are polyoxyethylene (20) sorbitan monolaurate, polyoxyethylene (20) sorbitan monopalmitate, polyoxyethylene (20) sorbitan monostearate, polyoxyethylene (20) sorbitan monooleate (Tween® 80), polyoxyethylene (20) sorbitan tristearate (Tween® 65), polyoxyethylene (20) sorbitan trioleate (Tween® 85), polyoxyethylene (4) sorbitan monostearate, polyoxyethylene (4) sorbitan monolaurate or polyoxyethylene (4) sorbitan monooleate.

Suitable polyalkoxylated glycerides are obtained for example by alkoxylation of natural or hydrogenated glycerides or by transesterification of natural or hydrogenated glycerides with polyalkylene glycols. Commercially available examples are polyoxyethylene glycerol ricinoleate 35, polyoxyethylene glycerol trihydroxystearate 40 (Cremophor® RH40, BASF AG) and polyalkoxylated glycerides like those obtainable under the proprietary names Gelucire® and Labrafil® from Gattefosse, e.g. Gelucire® 44/14 (lauroyl macrogol 32 glycerides prepared by transesterification of hydrogenated palm kernel oil with PEG 1500), Gelucire® 50/13 (stearoyl macrogol 32 glycerides, prepared by transesterification of hydrogenated palm oil with PEG 1500) or Labrafil M1944 CS (oleoyl macrogol 6 glycerides prepared by transesterification of apricot kernel oil with PEG 300).

A suitable fatty acid ester of polyalkylene glycols is, for example, PEG 660 hydroxystearic acid (polyglycol ester of 12-hydroxystearic acid (70 mol%) with 30 mol% ethylene glycol).

Suitable polyalkoxylated ethers of fatty alcohols are, for example, PEG (2) stearyl ether (Brij® 72), macrogol 6 cetylstearyl ether or macrogol 25 cetylstearyl ether.

In general, the tocopheryl compound corresponds to the formula below



wherein Z is a linking group, R¹ and R² are, independently of one another, hydrogen or C₁-C₄ alkyl and n is an integer from 5 to 100, preferably 10 to 50. Typically, Z is the residue of an aliphatic dibasic acid such as glutaric, succinic, or adipic acid. Preferably, both R¹ and R² are hydrogen.

5

It was found that solubilizers or combination of solubilizers having a defined HLB (hydrophilic lipophilic balance) value are preferred over other solubilizers.

10 The HLB system (Fiedler, H.B., Encyclopedia of Excipients, 5th ed., Aulendorf: ECV-Editio-Cantor-Verlag (2002)) attributes numeric values to surfactants, with lipophilic substances receiving lower HLB values and hydrophilic substances receiving higher HLB values.

15 Where a single solubilizer is employed it suitably has an HLB value of from 3.5 to 13, preferably from 4 to 11.

20 Where a combination of two or more pharmaceutically acceptable solubilizers is used the combination of pharmaceutically acceptable solubilizers suitably has an averaged HLB value in the range of from 4.5 to 12, preferably 5 to 11. The averaged HLB value may be computed by multiplying the HLB value of each individual solubilizer by the proportion of the individual solubilizer with regard to the total amount of solubilizers present and adding together the contributions of the individual solubilizers.

25 Quite unexpectedly, a combination of at least one solubilizer having a relatively high HLB value and at least one solubilizer having a relatively low HLB value proved particularly useful. The high HLB solubilizer suitably has an HLB value in the range of from 8 to 15, preferably 10 to 14. The low HLB solubilizer suitably has an HLB value in the range of from 3 to 6, preferably 3.5 to 5. The weight ratio of high HLB solubilizer and low HLB solubilizer may be in the range of from 9:1 to 1:9, preferably 5:1 to 1:5.

30

Solubilizers having an HLB value in the range of from 8 to 15 may be selected from Cremophor® RH40 (HLB 13), Tween® 65 (HLB 10.5), Tween® 85 (HLB 11). Preferred high HLB solubilizers are tocopheryl compounds having a polyalkylene glycol moiety.

35

The preferred tocopheryl compound is alpha tocopheryl polyethylene glycol succinate, which is commonly abbreviated as vitamin E TPGS. Vitamin E TPGS is a water-soluble form of natural-source vitamin E prepared by esterifying d-alpha-tocopheryl acid succinate with polyethylene glycol 1000. Vitamin E TPGS is available from Eastman Chemical Company, Kingsport, TN, USA and is listed in the US pharmacopoeia (NF).

40

Solubilizers having an HLB value in the range of from 3 to 6 may be selected from Span® (HLB 4.7), Span ® 80 (HLB 4.3), Labrafil M1944 CS (HLB 4.0) and Brij® 72 (HLB 4.9).

- 5 A preferred low HLB solubilizer is an alkylene glycol fatty acid monoester or a mixture of alkylene glycol fatty acid mono- and diester.

The preferred alkylene glycol fatty acid mono ester is a propylene glycol fatty acid mono ester, such as propylene glycol monolaurate (available under the trade name
10 LAUROGLYCOL® from Gattefossé, France). Commercially available propylene glycol lauric acid mono ester products consist of a mixture of mono- and dilaurate. Two propylene glycol monolaurate products are specified in the European Pharmacopoea (referenced "type I" and "type II" respectively). Both types are suitable for carrying out the present invention, with propylene glycol monolaurate "type I" being the most preferred.
15 This "type I" product having a HLB value of about 4 consists of a mixture having between 45 and up to 70% mono-laurate and between 30 and up to 55% of di-laurate. The "type II" product is specified according to Pharm. Eur. as having a minimum of 90% mono-laurate and a maximum of 10% of di-laurate.

- 20 Where a mixture of alkylene glycol fatty acid mono and diester is employed, this preferably contains at least 40% by weight of the mono ester, especially 45 to 95 % by weight, relative to the weight of the ester mixture.

Thus, in a preferred embodiment, the combination of solubilizers comprises (i) at least
25 one tocopheryl compound having a polyalkylene glycol moiety, preferably alpha tocopheryl polyethylene glycol succinate, and (ii) at least one alkylene glycol fatty acid monoester or a mixture of alkylene glycol fatty acid mono- and diester.

The pharmaceutically acceptable polymer may be selected from water-soluble polymers, water-dispersible polymers or water-swellaable polymers or any mixture thereof.
30 Polymers are considered water-soluble if they form a clear homogeneous solution in water. When dissolved at 20 °C in an aqueous solution at 2 % (w/v), the water-soluble polymer preferably has an apparent viscosity of 1 to 5000 mPa.s, more preferably of 1 to 700 mPa.s, and most preferably of 5 to 100 mPa.s. Water-dispersible polymers are
35 those that, when contacted with water, form colloidal dispersions rather than a clear solution. Upon contact with water or aqueous solutions, water-swellaable polymers typically form a rubbery gel.

Preferably, the pharmaceutically acceptable polymer employed in the invention has a
40 Tg of at least 40°C, preferably at least +50°C, most preferably from 80 ° to 180. °C. "Tg" means glass transition temperature. Methods for determining Tg values of the

- organic polymers are described in "Introduction to Physical Polymer Science", 2nd Edition by L.H. Sperling, published by John Wiley & Sons, Inc., 1992. The Tg value can be calculated as the weighted sum of the Tg values for homopolymers derived from each of the individual monomers, i , that make up the polymer: $T_g = \sum W_i X_i$ where W is the weight percent of monomer i in the organic polymer, and X is the Tg value for the homopolymer derived from monomer i . Tg values for the homopolymers may be taken from "Polymer Handbook", 2nd Edition by J. Brandrup and E.H. Immergut, Editors, published by John Wiley & Sons, Inc., 1975.
- Various additives contained in the solid dispersion product or even the active ingredient(s) itself may exert a plasticizing effect on the polymer and thus depress the Tg of the polymer such that the final solid dispersion product has a somewhat lower Tg than the starting polymer used for its preparation. In general, the final solid dispersion product has a Tg of 10 °C or higher, preferably 15 °C or higher, more preferably 20 °C or higher and most preferred 30 °C or higher.
- For example, preferred pharmaceutically acceptable polymers can be selected from the group comprising
- homopolymers and copolymers of N-vinyl lactams, especially homopolymers and copolymers of N-vinyl pyrrolidone, e.g. polyvinylpyrrolidone (PVP), copolymers of N-vinyl pyrrolidone and vinyl acetate or vinyl propionate,
 - cellulose esters and cellulose ethers, in particular methylcellulose and ethylcellulose, hydroxyalkylcelluloses, in particular hydroxypropylcellulose, hydroxyalkylalkylcelluloses, in particular hydroxypropylmethylcellulose, cellulose phthalates or succinates, in particular cellulose acetate phthalate and hydroxypropylmethylcellulose phthalate, hydroxypropylmethylcellulose succinate or hydroxypropylmethylcellulose acetate succinate;
 - high molecular polyalkylene oxides such as polyethylene oxide and polypropylene oxide and copolymers of ethylene oxide and propylene oxide,
 - polyvinyl alcohol-polyethylene glycol-graft copolymers (available as Kollicoat® IR from BASF AG, Ludwigshafen, Germany);
 - polyacrylates and polymethacrylates such as methacrylic acid/ethyl acrylate copolymers, methacrylic acid/methyl methacrylate copolymers, butyl methacrylate/2-dimethylaminoethyl methacrylate copolymers, poly(hydroxyalkyl acrylates), poly(hydroxyalkyl methacrylates),

polyacrylamides,

vinyl acetate polymers such as copolymers of vinyl acetate and crotonic acid, partially hydrolyzed polyvinyl acetate (also referred to as partially saponified "polyvinyl alcohol"),

polyvinyl alcohol,

oligo- and polysaccharides such as carrageenans, galactomannans and xanthan gum, or mixtures of one or more thereof.

Among these, homopolymers or copolymers of N-vinyl pyrrolidone, in particular a copolymer of N-vinyl pyrrolidone and vinyl acetate, are preferred. A particularly preferred polymer is a copolymer of 60 % by weight of the copolymer, N-vinyl pyrrolidone and 40 % by weight of the copolymer, vinyl acetate.

A further polymer which can be suitably used is Kollidon® SR (available from BASF AG, Ludwigshafen, Germany) which comprises a mixture of PVP and polyvinylacetate.

The solid dispersion product may be prepared by a variety of methods. The solid dispersion product may be prepared by a solvent evaporation method. In a solvent evaporation method, the at least one tyrosine kinase inhibitor, the at least one pharmaceutically acceptable polymer and the at least one pharmaceutically acceptable solubilizer are dissolved in a common solvent or combination of solvents and the solvents are removed from the solution obtained by evaporation.

Preferably, the solid dispersion product is prepared by melt-extrusion. The melt-extrusion process comprises the steps of preparing a homogeneous melt of the active ingredient or the combination of active ingredients, the pharmaceutically acceptable polymer and the solubilizers, and cooling the melt until it solidifies. "Melting" means a transition into a liquid or rubbery state in which it is possible for one component to become homogeneously embedded in the other. Typically, one component will melt and the other components will dissolve in the melt, thus forming a solution. Melting usually involves heating above the softening point of the pharmaceutically acceptable polymer. The preparation of the melt can take place in a variety of ways. The mixing of the components can take place before, during or after the formation of the melt. For example, the components can be mixed first and then melted or simultaneously mixed and melted. Usually, the melt is homogenized in order to disperse the active ingredients efficiently. Also, it may be convenient first to melt the pharmaceutically acceptable polymer and then to admix and homogenize the active ingredients.

Usually, the melt temperature is in the range of 70 to 250 °C, preferably 80 to 180 °C, most preferably 100 to 140 °C.

5 The active ingredients can be employed as such or as a solution or dispersion in a suitable solvent such as alcohols, aliphatic hydrocarbons or esters. Another solvent which can be used is liquid carbon dioxide. The solvent is removed, e.g. evaporated, upon preparation of the melt.

10 Various additives may be included in the melt, for example flow regulators such as colloidal silica; lubricants, bulking agents (fillers), disintegrants, plasticizers, stabilizers such as antioxidants, light stabilizers, radical scavengers, or stabilizers against microbial attack.

15 The melting and/or mixing takes place in an apparatus customary for this purpose. Particularly suitable are extruders or kneaders. Suitable extruders include single screw extruders, intermeshing screw extruders or else multiscrew extruders, preferably twin screw extruders, which can be corotating or counterrotating and, optionally, equipped with kneading disks or other screw elements for mixing or dispersing the melt. It will be appreciated that the working temperatures will also be determined by the kind of extruder or the kind of configuration within the extruder used. Part of the energy needed to melt, mix and dissolve the components in the extruder can be provided by heating elements. However, the friction and shearing of the material in the extruder may also provide a substantial amount of energy to the mixture and aid in the formation of a homogeneous melt of the components.

25 The extrudate exiting from the extruder ranges from pasty to viscous. Before allowing the extrudate to solidify, the extrudate may be directly shaped into virtually any desired shape. Shaping of the extrudate may be conveniently carried out by a calender with two counter-rotating rollers with mutually matching depressions on their surface. A broad range of tablet forms can be attained by using rollers with different forms of depressions. If the rollers do not have depressions on their surface, films can be obtained. Alternatively, the extrudate is moulded into the desired shape by injection-moulding. Alternatively, the extrudate is subjected to profile extrusion and cut into pieces, either before (hot-cut) or after solidification (cold-cut).

35 Additionally, foams can be formed if the extrudate contains a propellant such as a gas, e.g. carbon dioxide, or a volatile compound, e.g. a low molecular-weight hydrocarbon, or a compound that is thermally decomposable to a gas. The propellant is dissolved in the extrudate under the relatively high pressure conditions within the extruder and, 40 when the extrudate emerges from the extruder die, the pressure is suddenly released.

Thus the solvability of the propellant is decreased and/or the propellant vaporises so that a foam is formed.

- 5 Optionally, the resulting solid solution product is milled or ground to granules. The granules may then be filled into capsules or may be compacted. Compacting means a process whereby a powder mass comprising the granules is densified under high pressure in order to obtain a compact with low porosity, e.g. a tablet. Compression of the powder mass is usually done in a tablet press, more specifically in a steel die between two moving punches.
- 10 At least one additive selected from flow regulators, disintegrants, bulking agents (fillers) and lubricants is preferably used in compacting the granules. Disintegrants promote a rapid disintegration of the compact in the stomach and keep the liberated granules separate from one another. Suitable disintegrants are crosslinked polymers such as
- 15 crosslinked polyvinyl pyrrolidone and crosslinked sodium carboxymethyl cellulose. Suitable bulking agents (also referred to as "fillers") are selected from lactose, calcium hydrogenphosphate, microcrystalline cellulose (Avicel®), magnesium oxide, potato or corn starch, isomalt, polyvinyl alcohol.
- 20 Suitable flow regulators are selected from highly dispersed silica (Aerosil®), and animal or vegetable fats or waxes.
- A lubricant is preferably used in compacting the granules. Suitable lubricants are selected from polyethylene glycol (e.g., having a Mw of from 1000 to 6000), magnesium
- 25 and calcium stearates, sodium stearyl fumarate, talc, and the like.
- Various other additives may be used, for example dyes such as azo dyes, organic or inorganic pigments such as aluminium oxide or titanium dioxide, or dyes of natural origin; stabilizers such as antioxidants, light stabilizers, radical scavengers, or stabilizers
- 30 against microbial attack.
- Dosage forms according to the invention may be provided as dosage forms consisting of several layers, for example laminated or multilayer tablets. They can be in open or closed form. "Closed dosage forms" are those in which one layer is completely sur-
- 35 rounded by at least one other layer. Multilayer forms have the advantage that two active ingredients which are incompatible with one another can be processed, or that the release characteristics of the active ingredient(s) can be controlled. For example, it is possible to provide an initial dose by including an active ingredient in one of the outer layers, and a maintenance dose by including the active ingredient in the inner layer(s).
- 40 Multilayer tablets types may be produced by compressing two or more layers of granules. Alternatively, multilayer dosage forms may be produced by a process known as

"coextrusion". In essence, the process comprises the preparation of at least two different melt compositions as explained above, and passing these molten compositions into a joint coextrusion die. The shape of the coextrusion die depends on the required drug form. For example, dies with a plain die gap, called slot dies, and dies with an annular slit are suitable.

In order to facilitate the intake of such a dosage form by a mammal, it is advantageous to give the dosage form an appropriate shape. Large tablets that can be swallowed comfortably are therefore preferably elongated rather than round in shape.

A film coat on the tablet further contributes to the ease with which it can be swallowed. A film coat also improves taste and provides an elegant appearance. If desired, the film coat may be an enteric coat. The film coat usually includes a polymeric film-forming material such as hydroxypropyl methylcellulose, hydroxypropyl cellulose, and acrylate or methacrylate copolymers. Besides a film-forming polymer, the film coat may further comprise a plasticizer, e.g. polyethylene glycol, a surfactant, e.g. a Tween® type, and optionally a pigment, e.g. titanium dioxide or iron oxides. The film-coating may also comprise talc as anti-adhesive. The film coat usually accounts for less than about 5 % by weight of the dosage form.

The dosage forms of the invention are useful for treating proliferative disorders, especially tumors or cancers. The proliferative disorder may be selected from the group consisting of neurofibromatosis, tuberous sclerosis, hemangiomas and lymphangiogenesis, cervical, anal and oral cancers, eye or ocular cancer, stomach cancer, colon cancer, bladder cancer, rectal cancer, liver cancer, pancreas cancer, lung cancer, breast cancer, cervix uteri cancer, corpus uteri cancer, ovary cancer, prostate cancer, testis cancer, renal cancer, brain cancer, cancer of the central nervous system, head and neck cancer, throat cancer, skin melanoma, acute lymphocytic leukemia, acute myelogenous leukemia, Ewing's Sarcoma, Kaposi's Sarcoma, basal cell carcinoma and squamous cell carcinoma, small cell lung cancer, choriocarcinoma, rhabdomyosarcoma, angiosarcoma, hemangioendothelioma, Wilms Tumor, neuroblastoma, mouth/pharynx cancer, esophageal cancer, larynx cancer, lymphoma, multiple myeloma; cardiac hypertrophy, age-related macular degeneration and diabetic retinopathy.

The exact dose and frequency of administration depends on the particular condition being treated, the age, weight and general physical condition of the particular patient as well as other medication the individual may be taking, as is well known to those skilled in the art.

The following examples will serve to further illustrate the invention without limiting it.

Example 1: Preparation of solid dispersion products

Formulations of various compositions were produced as shown in Table 1 below. The
5 active ingredient (N-[4-(3-amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)-
urea ethanolate) was mixed in a turbula blender with a pre-granulated mixture of Kolli-
don VA64 (copolymer of 60 % by weight N-vinyl pyrrolidone and 40 % by weight vinyl
acetate) and the solubilizer(s). Additionally 1% of colloidal silicon dioxide was added to
improve flow properties. The powdery mixture was extruded in a Leistritz micro 18
10 GMP-extruder at the extrusion temperature and rotational speed as shown in table 1.

Table 1:

Example	ABT 869 wt %	Kollidon VA64 wt. %	Solubilizer 1 wt. %	Solubilizer 2 wt. %	T(°C)	U/min
A	7.5	87.5	Sorbitanmonolaurate 5	none	130-140	70
B	5	85	Sorbitanmonolaurate 10	none	140	150
C	5	90	Sorbitanmonolaurate 5	none	140	150
D	5	92	Sorbitanmonolaurate 3	none	140-145	150
E	5	90	Tween 80 5	none	140	150
F	5	90	TPGS* 5	none	140	150
G	5	90	Propylenglycol monolaurate** 5	none	140	150
H	5	85	Propylenglycol monolaurate** 6.2	TPGS* 3.8	140	150
I	5	90	Propylenglycol monolaurate** 3.1	TPGS* 1.9	140	150
K	5	89	Propylenglycol monolaurate** 4	TPGS* 2	140	150

Example	ABT 869 wt %	Kollidon VA64 wt. %	Solubilizer 1 wt. %	Solubilizer 2 wt. %	T(°C)	U/min
L	5	85	Sorbitanmonolaurate 10	4.81	140	150
M	5	90	Propylenglycol monolaurate** 3.1	TPGS* 1.9	140	150
N	5	80	Sorbitanmonolaurate 15	none	140	150
O	5	80	Sorbitanmonolaurate 15	none	125	150
P	5	80	Tween 80 15	none	140	150
Q	5	80	Tween 80 15	none	125	150
R	5	80	Propylenglycol monolaurate** 10	TPGS* 5	140	150
S	5	80	Propylenglycol monolaurate** 10	TPGS* 5	125	150

* tocopheryl polyethylene glycol 1000 succinate

** Type 1

Example 2: Bioavailability Evaluation

Protocol for the oral bioavailability studies

- 5 For bioavailability evaluation, extrudates as obtained in Example 1 were milled and filled into capsules. Each capsule contained 25 mg ABT 869.

The studies were run with liquid clinical formulation as reference (4.0 % by weight ABT 869 in ethanol-surfactant solution) in a two-treatment, two-period crossover study.

10

Dogs (beagle dogs, mixed sexes, weighing approximately 10 kg) received a balanced diet with 27 % fat and were permitted water ad libitum. Each dog received a 100 µg/kg subcutaneous dose of histamine approximately 30 minutes prior to dosing. A single dose corresponding to 25 mg ABT 869 was administered to each dog. The dose was followed by approximately 10 milliliters of water. Blood samples were obtained from each animal prior to dosing and 0.25, 0.5, 1.0, 1.5, 2, 3, 4, 6, 8, 10, 12 and 24 hours after drug administration. The plasma was separated from the red cells by centrifugation and frozen (-30 °C) until analysis. Concentrations of ABT 869 inhibitors were determined by reverse phase HPLC with low wavelength UV detection following liquid-liquid extraction of the plasma samples. The area under the curve (AUC) was calculated by the trapezoidal method over the time course of the study. Each dosage form was evaluated in a group containing 5-6 dogs; the values reported are averages for each group of dogs.

25

Table 2: Results of dog studies with a crossover study design

Example	C _{max} [µg/ml]	T _{max} [h]	Pt. Esti- mate C _{max} [*]	Pt. Esti- mate AUC [*]
N	0.77	1.1	0.83	0.82
G	0.51	1.0	1.04	1.11
I	0.46	1.4	0.93	1.07
K	0.56	1.9	0.68	0.8
R	0.84	1.1	1.04	1.04

*The values are reported as relative bioavailability-compared to the bioavailability of the liquid clinical formulation as reference.

5 Example 3: Manufacture of tablets

Following the procedure of example 1, an extrudate was obtained from the solid dispersion product ingredients listed in table 3 below. The extrudate was allowed to cool.
10 The solidified extrudate was milled and the powder was blended with the tableting excipients listed in table 3. A tablet press was used to prepare tablets containing 2.5 mg or 10 mg, respectively, of ABT-869.

Table 3: Tablet composition

15

Ingredient	% (w/w)
Solid dispersion product	
ABT-869 ethanolate	2.50
Kollidon VA64	39.75
Propylene glycol monolaurate (Type I)	5.00
Vitamin E-TPGS	2.50
Colloidal silicon dioxide, Type Aerosil 200	0.25
Tabletting excipients	
Mannitol	48.50
Colloidal silicon dioxide , Type Aerosil 200	1.00
Sodium stearyl fumarate	0.50

Example 4: Estimating Pharmacokinetics in humans

20 Tablets containing a 10 mg dose of ABT-869 ethanolate, as prepared above, were administered to 11 patients in the morning with 240 mL of water under fasting conditions (defined as no food or drink consumption except water and concomitant medications for 2 hours prior to dosing). Following dosing, 4-mL blood samples were collected for pharmacokinetic analyses at the following times: 0 (pre-dose), 1, 3, 4, 6, 8, 24 and 48
25 hours. These samples were analyzed for ABT-869 plasma concentrations using Liquid Chromatography/Tandem Mass Spectrometry (LCMS/MS). The lower limit of quantification (LLOQ) for the assay was 1.1 ng/mL.

Pharmacokinetic parameters including the maximum observed plasma concentration
30 (C_{\max}), time to C_{\max} (T_{\max}), the area under the plasma concentration-time curve (AUC) from 0 to time of the last measurable concentration (AUC_{0-48}) and AUC from 0 to infinite

time (AUC_{∞}) were determined by non-compartmental methods using WinNonlin Professional version 5.2 software.

- 5 AUC_{0-48} per mg of dose was $0.40 \pm 0.10 \mu\text{g} \cdot \text{h/mL/mg}$ (mean \pm SD) while AUC_{∞} per mg of dose was $0.55 \pm 0.17 \mu\text{g} \cdot \text{h/mL/mg}$. C_{max} per mg of dose was calculated to be $0.023 \pm 0.004 \mu\text{g/mL/mg}$. The ABT-869 tablet has a T_{max} of 2.8 ± 0.6 h. The intersubject variability in the ABT-869 tablet was 17% in C_{max} and 25% in AUC_{0-48} .

We claim:

1. A pharmaceutical dosage form which comprises a solid dispersion product of at least one tyrosine kinase inhibitor, at least one pharmaceutically acceptable polymer, and at least one pharmaceutically acceptable solubilizer.
2. The dosage form of claim 1, which upon contact with an aqueous liquid releases particles having an average particle size of less than about 1000 nm, the particles containing solubilised tyrosine kinase inhibitor.
3. The dosage form of claim 1, wherein the pharmaceutically acceptable solubilizer is selected from the group consisting of polyol fatty acid esters, polyalkoxylated polyol fatty acid esters, polyalkoxylated fatty alcohol ethers, tocopheryl compounds or mixtures of two or more thereof.
4. The dosage form of claim 1, wherein the pharmaceutically acceptable solubilizer has an HLB value in the range of from 3.5 to 13.
5. The dosage form of claim 1, comprising a combination of two or more pharmaceutically acceptable solubilizers.
6. The dosage form of claim 5, wherein the combination of pharmaceutically acceptable solubilizers has an averaged HLB value in the range of from 4.5 to 12.
7. The dosage form of claim 5, wherein the combination of pharmaceutically acceptable solubilizers comprises (i) at least one solubilizer having an HLB value in the range of from 8 to 15 and (ii) at least one solubilizer having an HLB value in the range of from 3 to 6.
8. The dosage form of claim 7, wherein the combination of pharmaceutically acceptable solubilizers comprises (i) at least one tocopheryl compound having a polyalkylene glycol moiety and (ii) at least one alkylene glycol fatty acid monoester or mixture of alkylene glycol fatty acid mono- and diester.
9. The dosage form of claim 8, wherein the tocopheryl compound is alpha tocopheryl polyethylene glycol succinate.
10. The dosage form of claim 8, wherein the alkylene glycol fatty acid monoester is propylene glycol monolaurate.
11. The dosage form of Claim 8, wherein the weight ratio of tocopheryl compound and alkylene glycol fatty acid ester is in the range of from 9:1 to 1:9.
12. The solid dosage form of claim 1, wherein said pharmaceutically acceptable polymer is a homopolymer or copolymer of N-vinyl pyrrolidone.

13. The solid dosage form of claim 1, wherein said pharmaceutically acceptable polymer is a copolymer of N-vinyl pyrrolidone and vinyl acetate.
- 5 14. The dosage form of claim 1, wherein said tyrosine kinase inhibitor is selected from the group consisting of sorafenib, dasatinib, lapatinib, imatinib, motesanib, vandetanib, MP-412, lestauritinib, XL647, XL999, tandutinib, PKC412, nilotinib, AEE788, OSI-930, OSI-817, sunitinib maleate, axitinib,
10 N-[4-(3-amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea (ABT869);
N-(4-(4-aminothieno[2,3-d]pyrimidin-5-yl)phenyl)-N'-(2-fluoro-5-(trifluoromethyl)-phenyl)urea; or salts or hydrates or solvates thereof,
or combinations thereof.
- 15 15. The dosage form of claim 1, wherein said tyrosine kinase inhibitor is poorly water-soluble.
16. The dosage form of claim 1, wherein said tyrosine kinase inhibitor comprises at
20 least one urea moiety in its molecular structure.
17. The solid dosage form of claim 1, containing at least one additive selected from flow regulators, disintegrants, bulking agents and lubricants.
- 25 18. The dosage form of claim 1, wherein the solid dispersion product comprises from about 0.5 to 40 % by weight of said at least one tyrosine kinase inhibitor, 40 to 97.5 % by weight of said at least one pharmaceutically acceptable polymer, 2 to 20 % by weight of said at least one solubilizer, and 0 to 15 % by weight of additives.
- 30 19. The dosage form of claim 1 wherein the solid dispersion product is a melt-processed, solidified mixture.
- 35 20. The dosage form of claim 1 wherein the solid dispersion product is obtained by dissolving the at least one tyrosine kinase inhibitor, the at least one pharmaceutically acceptable polymer and the at least one pharmaceutically acceptable solubilizer in a common solvent or combination of solvents and evaporating the solution obtained.
- 40 21. The dosage form of claim 1 wherein said tyrosine kinase inhibitor is N-[4-(3-amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea (ABT 869), the dosage form, when administered to a human patient, producing a plasma profile characterized by a C_{max} for ABT 869 from about 0.015 $\mu\text{g/mL/mg}$ to about 0.027 $\mu\text{g/mL/mg}$ after a single dose.
- 45

22. The dosage form of claim 1 wherein said tyrosine kinase inhibitor is N-[4-(3-amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea (ABT 869), the dosage form, when administered to a human patient, producing a plasma profile characterized by a T_{max} for ABT 869 from 1 to about 3 hours after a single dose.
23. The dosage form of claim 1 wherein said tyrosine kinase inhibitor is N-[4-(3-amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea (ABT 869), the dosage form, when administered to a human patient, producing a plasma profile characterized by a AUC_{0-48} per mg of ABT 869 from about $0.23 \mu\text{g}\cdot\text{hr}/\text{mL}/\text{mg}$ to about $0.56 \mu\text{g}\cdot\text{hr}/\text{mL}/\text{mg}$ per mg of dose after a single dose.
24. The dosage form of claim 1 wherein said tyrosine kinase inhibitor is N-[4-(3-amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea (ABT 869), the dosage form, when administered to a human patient, producing a plasma profile characterized by a $AUC_{0-\infty}$ per mg of ABT 869 from about $0.27 \mu\text{g}\cdot\text{hr}/\text{mL}/\text{mg}$ to about $0.81 \mu\text{g}\cdot\text{hr}/\text{mL}/\text{mg}$ per mg of dose after a single dose.
25. A method of treating proliferative disorders, comprising administering the dosage form of claim 1 to a subject in need thereof.
26. The method of claim 25, wherein the proliferative disorder is selected from tumors or cancers.
27. The method of claim 25, wherein the proliferative disorder is selected from the group consisting of neurofibromatosis, tuberous sclerosis, hemangiomas and lymphangiogenesis, cervical, anal and oral cancers, eye or ocular cancer, stomach cancer, colon cancer, bladder cancer, rectal cancer, liver cancer, pancreas cancer, lung cancer, breast cancer, cervix uteri cancer, corpus uteri cancer, ovary cancer, prostate cancer, testis cancer, renal cancer, brain cancer, cancer of the central nervous system, head and neck cancer, throat cancer, skin melanoma, acute lymphocytic leukemia, acute myelogenous leukemia, Ewing's Sarcoma, Kaposi's Sarcoma, basal cell carcinoma and squamous cell carcinoma, small cell lung cancer, choriocarcinoma, rhabdomyosarcoma, angiosarcoma, hemangioendothelioma, Wilms Tumor, neuroblastoma, mouth/pharynx cancer, esophageal cancer, larynx cancer, lymphoma, multiple myeloma; cardiac hypertrophy, age-related macular degeneration and diabetic retinopathy.
28. A method of preparing a solid dosage form of claim 1 which comprises:
- preparing a homogeneous melt of said at least one tyrosine kinase inhibitor, said at least one pharmaceutically acceptable polymer and said at least one solubilizer, and
 - allowing the melt to solidify to obtain a solid dispersion product.

29. The method of claim 28, additionally comprising grinding said solid dispersion product and compressing said solid dispersion product into a tablet.
- 5 30. The method of claim 28, additionally comprising grinding said solid dispersion product and filling said solid dispersion product into a capsule shell.
31. The method of claim 28, wherein the melt is shaped into a film or a foam before being allowed to solidify.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2007/062101

A. CLASSIFICATION OF SUBJECT MATTER
INV. A61K9/20 A61K31/505

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, EMBASE, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99/03854 A (NOVARTIS AG ET AL.) 28 January 1999 (1999-01-28) page 17, lines 23-29; claims 11,12; example 4 -----	1-31
X	WO 2005/011647 A (ASTRA ZENECA AB ET AL.) 10 February 2005 (2005-02-10) page 38, lines 15-28 page 39, lines 12-26; claims 1-19 -----	1-31
X	US 2006/078617 A1 (F. SCHUECKLER) 13 April 2006 (2006-04-13) claims 1-33; example 4 -----	1-31
X	US 2006/154941 A1 (LE HUANG) 13 July 2006 (2006-07-13) claims 1-19; table 1 -----	1-31
	-/--	

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

17 January 2008

Date of mailing of the international search report

25/01/2008

Name and mailing address of the ISA/
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

GLIKMAN, J

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2007/062101

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2006/178387 A1 (K. FUJIMOTO-OUCHI ET AL.) 10 August 2006 (2006-08-10) paragraphs [0093], [0112] -----	1-31
A	DATABASE CA CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; XP002464017 retrieved from STN Database accession no. 146: 32918 abstract & CN 1 857 204 A (QINGLUN KONG) 8 November 2006 (2006-11-08) -----	1-31

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2007/062101

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 25-27 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers allsearchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search reportcovers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2007/062101

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9903854	A	28-01-1999	AT 251152 T	15-10-2003
			AU 740713 B2	15-11-2001
			AU 8975998 A	10-02-1999
			BR 9810920 A	15-08-2000
			CA 2296604 A1	28-01-1999
			CN 1264375 A	23-08-2000
			CZ 20000149 A3	17-05-2000
			DE 69818674 D1	06-11-2003
			DE 69818674 T2	12-08-2004
			DK 998473 T3	02-02-2004
			EP 0998473 A1	10-05-2000
			ES 2209194 T3	16-06-2004
			HK 1028599 A1	30-04-2004
			HU 0003230 A2	28-06-2001
			ID 24093 A	06-07-2000
			JP 3276359 B2	22-04-2002
			JP 2001510192 T	31-07-2001
			NO 20000227 A	17-01-2000
			NZ 502295 A	21-12-2001
			PL 338129 A1	25-09-2000
			PT 998473 T	27-02-2004
			RU 2208012 C2	10-07-2003
			SK 432000 A3	12-06-2000
			TR 200000060 T2	21-09-2000
			TW 491845 B	21-06-2002
			US 6894051 B1	17-05-2005
			ZA 9806362 A	22-01-1999

WO 2005011647	A	10-02-2005	EP 1651233 A2	03-05-2006
			JP 2007500176 T	11-01-2007
			US 2007141141 A1	21-06-2007

US 2006078617	A1	13-04-2006	NONE	

US 2006154941	A1	13-07-2006	CN 101016266 A	15-08-2007

US 2006178387	A1	10-08-2006	NONE	

CN 1857204	A	08-11-2006	NONE	
